

Technical Support

Products #7706 & #7707

De-Stress™

For some mammalian species (including humans), the newborn is an external embryo. The milk they receive becomes the umbilical cord which links the newborn to the maternal organism (the mother). A baby's enzymatic system is different from that of the adult, especially concerning low pepsin levels, thereby displaying predominant trypsic activity.

Observing the baby's calm state after drinking warm milk, a research team from Nancy University observed the link between the trypsic activity in the milk and the calm state of the baby. This led to research on the milk casein hydrolysate fractions and peptides with anxiolytic activity.

DeStress™, Defined

DeStress™ is a patented, specially produced milk protein hydrolysate, which constitutes a specific peptide concentrate, having demonstrated anxiolytic activity. Anxiolytic activity refers to the action of medications prescribed for anxiety symptoms.

The Peptide


Identification of the peptide began with the laboratory production of an alpha-S1 casein trypsic hydrolysate (α_{s1} -casein), which demonstrated anxiolytic activity both *in vitro* (GABA_A receptor-binding test and peripheral-type benzodiazepine binding assays)¹ and *in vivo* (elevated plus maze test and conditioned defensive burying test, in rats).^{1,2}

Amino acid sequencing identified the peptide as a "benzodiazepine-like decapeptide"¹ Utilizing molecular separation techniques, the peptide demonstrating specific anxiolytic activity, was isolated from the trypsic hydrolysate.

The results of circular dichroism, two-dimensional NMR spectroscopy, and molecular modeling characterized the peptide as a 3₁₀ helix structure, initiated and terminated by an alpha-turn. In addition to the structural determination, it was observed that the decapeptide exhibited an amphiphilic characteristic, with the hydrophobic side chains located on one side of the molecule, and the hydrophilic side chains on the opposite side, thus indicating that it was "essentially flexible"³. Accordingly, the peptide was hypothesized to have the ability to cross both hydrophobic and hydrophilic membranes. Further classification of the peptide structure was confirmed by three-dimensional molecular modeling.⁴

Pre-Clinical Studies

The action of DeStress™ casein hydrolysate was demonstrated in randomized double blind studies, utilizing Wistar rats. The anxiolytic activity, reported as being similar to diazepam, was confirmed in these subjects. Results indicated that at a dosage of 3mg/kg, various activities demonstrated significance, including a decreased duration of probe burying ($P<0.005$), head stretches toward the probe ($P<0.01$), and the percent approaches towards the probe followed by retreats ($P<0.01$).¹ Additionally, rats given a 3mg/kg dose of the α_{s1} -casein tryptic hydrolysate (DeStress™) were demonstrated to show a significant reduction in the action of pentylenetetrazole (PTZ) ($P<0.002$), which is similar to the action of diazepam (2mg/kg).¹ The DeStress™ casein hydrolysate was also shown to have a "protective effect" on sleep, when subjects were exposed to chronic mild stress, observed as a



"maintenance of slow wave sleep duration," along with a minor increase in paradoxical sleep duration.⁵

Human Clinical Trials

In a double-blind, randomized, crossover, placebo-controlled trial, sixty-three female volunteers were randomly assigned to receive either α_{s1} -casein (150mg/day) or placebo. Following a thirty-day treatment program, the α_{s1} -casein hydrolysate group was observed to have reduced stress-related symptoms. The study correlated a reduction in stress related symptomatology to improvements in the following areas; digestion ($P<0.05$), cardiovascular ($P<0.05$), intellectual ($P<0.05$), emotional ($P<0.05$), and social ($P<0.05$) complications.⁶

In a subsequent study forty-two healthy male volunteers partook in a randomized, placebo controlled, double blind trial. The results indicated an "antistress activity in human subjects", as evidenced by a significant decrease in plasma cortisol in the α_{s1} -casein group (20.69%) ($t=3.73$; $P=0.001$), compared to placebo (3.39%) ($t=1.05$; $P=0.30$). Additionally, both the systolic and diastolic blood pressures were significantly



higher in the control group as compared to the α_{s1} -casein group. A decrease in heart rate ($P=0.05$) was also noted in the α_{s1} -casein group, compared to placebo.

In another double-blind crossover study, sixty-three women volunteers, presenting with at least one symptom of stress, were randomized to either receive DeStress™ or placebo for thirty days. A 30d washout period was then performed, followed by a reversal in the treatment regiments. The subjects answered questionnaires on the 1st, 15th, and 30th day of each treatment period. The questionnaires covered three main areas and eight sub-areas potentially affected by stress. The results indicated a significant improvement on the digestive system (58% vs. 38.6% for placebo) and the cardiovascular system (53.4% vs. 21.8% for placebo) in those subjects initially demonstrating the highest intensities for their major symptoms. Additionally, a significant improvement in five areas was noted following the 30d treatment with DeStress™ in subjects initially demonstrating the highest intensities of symptoms (>4/day). The areas of improvement were noted in digestive function (66.1% vs. 36% for placebo), cardiovascular problems (48% vs. 35.5% for placebo), intellectual disorders (64.8 vs. 36.7% for placebo), emotional disorders (43.8% vs. 23.5% for placebo) and social problems (36.7% vs. 22.5% for placebo).⁸

Thus confirming evidence indicates that DeStress™ has anxiolytic activity. As such administration has shown beneficial results towards alleviating symptoms of stress, including digestive, cardiovascular and anxiety effects.

References

1. Miclo L, Perrin E, Driou A, Papadopoulos V, Boujard N, Vanderesse R, Boudier JF, Desor Didier, Linden G, and Gaillard, JL. Characterization of -casozepine, a tryptic peptide from bovine α_{s1} -casein with benzodiazepine-like activity. *FASEB J.* 2001 Aug;15(10):1780-2.
2. Violle N, Messaoudi M, Lefranc-Millot C, Desor D, Nejdi A, Demagny B, Schroeder H. Ethological comparison of the effects of a bovine α_{s1} -casein tryptic hydrolysate and diazepam on the behaviour of rats in two models of anxiety. *Pharmacology, Biochemistry and Behavior* 84 (2006) 517-523.
3. Lecouvey M, Frochot C, Miclo L, Orlewski P, Driou A, Linden G, Gaillard JL, Marraud M, Cung MT, Vanderesse R. Two-dimensional 1H-NMR and CD structural analysis in a micellar medium of a bovine α_{s1} -casein fragment having benzodiazepine-like properties. *Eur J Biochem.* 1997. 248:872-878.
4. Kumosinski TF, Brown EM and Farrell HM. Three-dimensional Molecular Modeling of Bovine Caseins: α_{s1} -casein. *1991 Sep;74(9):2889-95.*
5. Guesdon B, Messaoudi M, Lefranc-Millot C, Fromentin G, Tome D, Even PC. A tryptic hydrolysate from bovine milk alpha α_{s1} -casein improves sleep in rats subjected to chronic stress. *Peptides.* 2006 Jun;24(6):1476-82.
6. Kim JH, Desor D, Kim YT, Yoon WJ, Kim KS, Jun JS, Pyun KH, Shim I. Efficacy of α_{s1} -casein hydrolysate on stress-related symptoms in women. *Eur J Clin Nutr.* 2007 Apr;61(4):536-41.
7. Bourdon, L., Messaoudi, M., Desor, D., Lefranc-Millot, C., Demagny, B. Effects of a tryptic hydrolysate from bovine milk α_{s1} -casein on hemodynamic responses in healthy human volunteers facing successive mental and physical stress situations. 2005. *Eur J Nutr.* 44(2):128-132.
8. Santure, M. Anti-Stress Efficacy of De-Stress™ on Stressed Women. *Nutra Foods, Volume 4, No. 1, 63-65 (2005).*

Product Information

DeStress™ is available in bottles of 10 and 30 capsules.

Supplement Facts

Serving Size: 1 Capsule	Amount Per Serving	% Daily Value
De-Stress™ hydrolysate † (as hydrolyzed casein concentrate)	150 mg	*
*Daily Value not established		

Other ingredients: Gelatin, water and glycerin.

Contains ingredients derived from milk.

† De-Stress™ supplies a decapeptide, as determined by its amino acid sequence, having anxiolytic activity. This statement has not been evaluated by the U.S. Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Patent # 5,846,939

RECOMMENDATION: One (1) capsule as a dietary supplement or as otherwise directed by a healthcare professional. De-Stress™ can be taken during the day at time of intense stress or before bedtime.

KEEP OUT OF REACH OF CHILDREN
Store in a cool, dry area.
Sealed with an imprinted safety seal for your protection.

NDC #55146-07706 (10 Capsules)
NDC #55146-07707 (30 Capsules)

For additional information call: (800) 373-1373
Metabolic Management
P.O. Box 546
Grant Park IL 60940
support@metabolicmanagement.com

